

A new species of *cucullanus* (Nematoda: Cucullanidae) from marine edible fish *Otolithus ruber* (Sciaenidae) based on light and Scanning Electron Microscopy.

Yasmin Akhtar^{1} and Bilqees Fatima Mujib¹*

¹Department of Biological Sciences Jinnah University for Women 5- C Nazimabad Karachi- 4600

ABSTRACT

Based on light and scanning electron microscopical observations, a new species of *Cucullanus* Müller, 1777 (Nematoda: Cucullanidae) is described from the intestine of marine edible fishes *Otolithus ruber* (Schneider, 1801), were collected (February 2006 to July 2007) from fresh landing of Karachi coast, Pakistan. Detailed light and scanning electron microscopy revealed some important taxonomical features like, Broader anteriorly, posterior end pointed, provided with small bifurcated spike at the tip of the tail. Lateral alae absent. Oral opening dorsoventrally elongate, surrounded by narrow membranous flange or collarete supported by row of numerous teeth. Three submedian cephalic papillae and a pair of prominent lateral amphids are present. Pseudobuccal capsule or esophastome wider than posterior part of esophagus. Deirids simple, just anterior to esophagus and intestinal junction. Tail broadly conical, bifurcated at the tip.

Key words: Parasitic nematode; new species; *Otolithus ruber*; fish; Intestine, Karachi coast; Pakistan.

INTRODUCTION

During a survey of nematodes from fishes of Karachi coast, Pakistan, a new species of the nematode of the family Cucullanidae Cobbold 1864 from the intestine of *Otolithus ruber* (Sciaenidae) was recovered, during the year 2006 to 2007. This species is described here in detail and is regarded a new species.

MATERIALS AND METHODS

Nematodes collected from the intestine of *Otolithus ruber* (2 male and 4 female specimens) were processed for light and electron microscopy.

Examination of specimens for nematode parasites

A total of 20 fish specimens were subjected to parasitological examination. The body cavity of fish was opened, and the gut and liver were removed by

cuts in the region of the anus and the division between esophagus and anterior stomach. The section of the gut in a large Petri dish was opened with a longitudinal cut, and the whole inner surface lightly scraped to remove the parasites with mucus. The nematode recovered were washed in physiological saline and preserved in 70% ethanol. For light microscopy the nematodes were cleared in glycerin. Diagrams were prepared with a camera Lucida E 200 Nikon Drawing Tube. Measurements are given length by width in millimeters. Specimens for scanning electron microscopy were fixed in cold 4% glutaraldehyde in buffer (pH=7.2) and kept in it for 24 hours, then dehydrated through a graded series of alcohols, infiltrated with amyl acetate, after critical point drying mounted on stubs, coated with gold and photographs were taken with the help of SEM. Joel Japan JSM 6380A at an accelerating voltage of 15KV at Karachi University, central laboratory. The SEM measurements are in micrometers.

*Corresponding author: yasminhameedi1@yahoo.com

RESULTS

A total of 20 fish specimens were subjected to parasitological examination. From three infected fish specimens, a total of six nematodes including (2 male and 4 female) were collected. These nematodes were recovered from the intestine. The new nematode species was identified and compared with the available literature (Yamaguti, 1961; Sood, 1989; Bilqees *et al.* 2005; Akhtar & Bilqees, 2006 and Bilqees, 2007).

Taxonomic summary:

Cucullanus aliyai n. sp.
(Fig.1-2)

Order: Spiruridea Diesing 1861
Family: Cucullanidae Cobbold 1864
Sub family: Cucullaninae Yorke & Maplestone
1926
Genus: Cucullanus Muller, 1777

Type host: *Otolithus ruber*(Schneider, 1801)
Site of infection: Intestine
Type locality: Fish harbor, Karachi coast Pakistan
Prevalence: 15 % (20 fish examined / 3 fish
infected)
Intensity: 2 (with 2 male and 4 females).
Holotype(male): JUW. N. 40
Allotype(female): JUW. N. 41

Male: (2 specimens including holotype): Length of body 2.02-2.29, greatest width 0.35-0.40, entire esophagus 0.36-0.38 in length. Its minimum width is in the middle 0.10, Length of esophastome is 0.18-0.20, maximum width at esophastome 0.13-0.15, posterior club- shaped part of esophagus in 0.34-0.37 in length and 0.09-0.11 in width .Nerve ring at a distance of 0.22-0.29 from the anterior extremity. Testis situated in posterior half of body or slightly anteriorly. Precloacal sucker is absent but few weak muscle fibers can be send. Spicules sub equal and 0.59-0.61 and 0.70-0.77 in length. Gubernaculum is absent. 12 pairs of caudal papillae are present including, 7 pairs preanal, 4 pairs are post anal and




Fig.1. (A-E) *Cucullanus aliyai* n.sp.
A. Anterior end of male (holotype)
B. Psterior end of male
C. Head of male
D. Anterior end of female
E. Posterior end of female

Female: (4 specimens including allotype): Length of body is 3.24, maximum width is 0.30. Entire esophagus is 0.50 in length, Greatest width 0.15 at esophastome , minimum width almost at the middle. Posterior club-shaped portion 0.15 in widths. Distance of nerve ring from the anterior extremity 0.30. Vulva 0.17-0.18 post equatorial, prominent, vulvar lips elevated at a distance of 1.80 from the anterior extremity. each containing elongated gland cells, extruding out of body wall. Vagina long, muscular tube and is directed upward. Uterus thin- walled,




Figure 2. *Cucullanus aliyaii* n.sp. allotype

- A. Cephalic end(Enface View)
- B. Anterior region showing head and deirids
- C. Tail region (showing bifurcation)
- D. Anal Region

containing numerous relatively large, thin- shelled eggs 0.07x0.04. Tail is 0.011 in length.

REMARKS

The genus *Cucullanus* Muller, 1777(Cucullanidae) contains a large number of species parasitizing various fresh water and marine water fishes around the world. Due to minute differences in the morphology, a detail comparison among species is very complicated. Therefore the present new species *C. aliyaii* is compared with the species recovered from South Asia, especially Pakistan and some related species from other parts of the world. The species reported from marine fishes of Pakistan are all from fishes of Karachi coast. These include *C.armatus* (Yamaguti, 1954) reported by Rasheed (1968) from *Tachysurus serratus*; *C. diminutus* (Rasheed, 1968) from *Stromateus niger*. Rasheed (1968) also reported

C.hians (Dujardin,1945) from *Lates cacarifier*, *Sciaena* sp., *Belone strongylurus*, *Acanthopagrus berda* and *Sparus* sp. At the same time Rasheed described *C. identatus* from *Polynemus tetradactylus* and *Pristopoma hasta*. He also redescribed *C fastigatus* (Chandler, 1935) from *Pristopoma hasta*. *C. theraponi* (Rasheed, 1968) was described from *Hilsa* sp., and *Therapon* sp., *C. bilqeesi* (Petter, 1974) was originally described by Bilqees *et al.*, (1971) as *C.elongatus* from *Erethistes elongate* and Petter renamed it as the species name was pre-occupied. *C. quadrii* (Bilqees, & Fatima,1980) from *Arius serratus*; *C. olivaceus* (Akram, 1975-1976) was described from *Tachysurus serratus* and *Pomadosys olivaceus*. Akram (1975-1976) also described *C. sparus* from *Sparus spinifer* and *Arius dussumieri* and *C. tachysuri* from *Tachysurus serratus* and *T. dussumieri*. *Indocucullanus karachii* (Zaidi & Khan, 1975) from the fish *Engraulis indica* was transferred to the genus *Cucullanus* by Soota (1983) with a combination as *C. karachii*.

Petter (1974) regarded *C. tachysurusi* as *C. arabiense* (Ali & Kalyankar, 1967) from Maharashtra, Mangalore and Karnataka. Petter (1974) restricted its occurrence in Siluriform hosts and gave only three pair of preanal caudal papillae for the species.

Yamaguti, (1961) has listed 60 species in the genus *Cucullanus* including the genotype. In addition to this Gupta and Masoodi (1982) described *C. sootai* and listed another 30 species including *C. ritali* (Karve, 1952); *C. jaiswali* (Ali, 1956); *C.indica* (Agrawal, 1965); *C. pseudotropi* (Agrawal,1967); *C. arabianse* (Ali and Kalyankar,1967; Petter, 1974); *C. theraponi* (Rasheed,1968); *C.pangasius* (Soota and chaturvedi., 1971); *C. jalnaensis*; *C. alii* Kalyankar, (1971); *C. malvanae* Kalyankar, (1971); *C. tachysuri* (Kalyankar, 1971); *C. bilqeesi*, (Bilqees *et al.*, 1971; Petter, 1974); *C. carioca* (Vicente and Fernandes, 1973); *C. rougetae* (Vicente and Dos, 1974); *C. bagrae* (Petter,1974); *C. karachii* (Zaidi & Khan, 1975); *C.olivaceus* (Akram, 1975-1976); *C. guerrori* (Arya and Jhonson, 1975); *C. ariusi* (Srivastava and Gupta, 1976); *C. sciaenai* (Gupta and Gupta, 1979); *C. rivulatus* (Soota and Sarkar,

1980); *C. trichiurisi* (Gupta and Naqvi, 1983); *C. simhai* (Gupta and Naqvi, 1983); *C. thapari* (Gupta and Srivastava, 1984) and *C. mastacembeli* (Gupta and Srivastava, 1984).

Later on more species have been described both from fresh water and marine fishes of various localities including *C. (Truttaedacnitis) truttae* (Moravec, 1979); *C. campanae* (Labre and Petter, 1984); *C. fugianensis* (Wang, 1984); *C. brevispiculus* (Moravec *et al.*, 1993); *C. riograndensis* (Fortes *et al.*, 1993a); *C. fabrigasi* (Fortes *et al.*, 1993b); *C. mexicanus* (Capseta-Mandujanoi, 2000); *C. marplatensis* (Ingrid *et al.*, 2002); *C. oceaniensis* (Moravec *et al.*, 2005); *C. pedroi* (Timi and Lanfranchi, 2006). David *et al.*, (2007) describe *C. pargi* from the grey snapper *Lutjanus griseus* of the southern coast of Quintana Roo, Mexico. Moravec *et al.*, (2008) also describe *C. maldivensis* from marine fishes of Maldiv Islands.

The present species *C. aliyaii* is different from *C. Alti* (Kalyankar, 1971) Peter, 1974; *C. arabianse* (Ali & Kalyankar, 1967; Petter, 1974); *C. ariusi* (Srivastava & Gupta, 1976) Soota, (1983); *C. armatus* (Yamaguti, 1954); *C. bulbosa* (Lane, 1916) Barreto, 1918; *C. chrysophrydis* (Gendre, 1927); *C. guerreroi* (Arya & Johnson, 1975) Soota, 1983; *C. jalnaensis* (Kalyankar, 1971) Petter, 1974; *C. karachii* (Zaidi & Khan, 1975) Soota, 1983; *C. malvanae* (Kalyankar, 1971) Petter, 1974; *C. pangasius* (Soota & Chaturvedi, 1971); *C. ritai* Karve, 1952; *C. rivulatus* Soota & Sarker, 1980; *C. sciaenai* (Gupta & Gupta, 1979) Soota, 1983; *C. theraponi* (Rasheed, 1968); *C. indica* (Agrawal, 1965); *C. tachysuri* (Kalyankar, 1971) Petter, 1974; *C. mystusi* (Gupta & Naqvi, 1985) Gupta & Masoodi, 1986; *C. vinodae* (Gupta & Naqvi, 1985) Gupta & Masoodi, 1986; *C. schizothoraxi* Gupta & Masoodi, 1986; *C. sootai* (Gupta & Masoodi, 1986); *C. fujianensis* (Wang, 1984); *C. baylisi* (Lakshmi, 2000); as all these species have a gubernaculum. While gubernaculum is absent in the present new species. Species described by Moravec *et al.*, (1993a-b, 1997) also have gubernaculum. Similarly gubernaculum is present in *C. riograndensis* (Fortes *et al.*, 1993a) and *C.*

fabrigasi (Fortes *et al.*, 1993b).

Present species *C. aliyaii* is also different from *C. marplatensis* (Ingrid *et al.*, 2002); *C. bonaerensis* (Lafranchi *et al.*, 2004); *C. oceaniensis* (Moravec *et al.*, 2005); *C. pedroi* (Timi and Lanfranchi, 2006); *C. pargi* (David *et al.*, 2007) and from *C. maldivensis* (Moravec *et al.*, 2008) in variable combination of characters, specially in size of esophagus, spicule length and number and position of caudal papillae in male. But the present species *C. aliyaii* resemble in the presence of deirids with *C. oceaniensis* (Moravec *et al.*, 2005). *C. bilqeese* (Bilqeese *et al.*, 1971) Petter, 1974 and *C. olivaceus* are described only by females. These are also different from the present species in the size of body, esophagus and eggs. The presence of deirids are not reported previously from Pakistan. As mentioned the morphological characters and presence of deirids, the present nematodes of the genus *Cucullanus* appear an undescribed species for which the name *C. aliyaii* n. sp., is proposed.

Soota (1983) regarded *C. quadrii* as synonym of *C. armatus*. This species was originally described by Yamaguti (1954) from *Tachysurus* sp., (*Arius* sp.) in Borneo and later on by Rasheed (1968) from *T. serratus*. Even if *C. quadrii* is considered synonym to *C. armatus*, the present species is different having different host, lacking a gubernaculum and preloacal sucker, longer subequal spicule (0.59-0.61 and 0.70-0.77) in length), vulvular lips are prominent elevated and protruded. The caudal papillae in the present species are 12 in number and in *T. armatus* caudal papillae are (9-10) pairs. Characters such as the presence of deirids, absence of gubernaculum, preloacal sucker, number of caudal papillae, spicules length, and morphology of vulva, egg sizes and tail length are considered valuable differentiating diagnostic features. Therefore, present specimens are regarded a new species and its name *C. aliyaii* n.sp. is proposed. Present species resemble in some morphological features with *C. pakistanensis* Bilqeese *et al.*, 2005, including sub equal spicules, length of tail and bifurcation at its tip, absence of gubernaculum and preloacal sucker.

But body size, length of esophagus, presence of deirids and morphology of vulva is different from *C. pakistanensis*.

The species in which gubernaculum is absent such as *C. bengalensis* (Gupta and Masoodi, 1985), But the caudal papillae are same (12 in numbers) as in the present species.

Gubernaculum is present in *C. armatus* (Yamaguti, 1954); *C. bulbosa* (Barreto, 1918); *C. indica* (Agrawal, 1965); *C. tachysuri* (Kalyankar, 1971) Petter, (1974); *C. mystusi*, (Gupta and Masoodi, 1986); *C. riograndensis*, (Fortes *et al*; 1993a) *C. fabrigasi* (Fortes *et al*; 1993b); *C. baylisi* (Lakshmi, 2000). Lakshmi (2000) described *C. baylisi* from the related host *Arius thalassinus* in India. Present new species is also separated from *C baylisi* which has a gubernaculum and much smaller spicules (0.237- 0.198mm in length), smaller eggs (0.069 x 0.048) and smaller tail (0.348 in female and 0.16-0.18in male). in addition a large number of sessile caudal papillae are also present in *C. baylisi*. Lakshmi probably did not consider *C. quadrii* a synonym of *C. armatus* or was unaware about this.

REFERENCES

Agarwal, V. 1965. Some new nematode parasites from fresh water fishes of Lucknow Indian. J. Helminth., 17: 1-17.

Akram, M. 1975-1976. A new nematode from the marine fish of Karachi coast. Sind Univ. Res. Jour. (Sci. Ser.),9: 89-91.

Akhtar, Y. and Bilqees, F. M. 2006. Checklist of nematodes of marine fishes of Pakistan. Pak. J. Nematol; 24(20): 205-216.

Ali, S.M. 1956. Studies of the nematode parasites of fishes and birds found in Hyderabad. Indian. J. Helminth. 8(1956): 1-83.

Ali, S.M. and Kalyankar, S.D. 1967. *Indocucullanus arabianse* n.sp., from the intestine of *Tachysurus*

maculates (cat fish) in India. Indian. J. Helminth., 18, Seminar Suppl. (1966): 74-76.

- Arya, S. N. and Johnson, s. 1975. A new Cucullanoid nematode from *Cybium guttatum* from Indian water(Spiruroidae, Cucullanidae) M. Sociedad de Ciencias Naturales La Salle, 35(102): 291-295.
- Barreto, A.L. de Barros, 1918. Notas helminthologicas 111. *Cucullanus pulcherrimus*. Brazil Med., 32: 137-138.
- Barreto, A.L. 1922. Revisao da familia Cucullanidae Barreto, 1916. Mems. Inst. Oswaldo Cruz., 14: 68-87.
- Baylis, H. A. 1932. A new nematode of the genus *Cucullanus* from New Zealand. Ann. And Mag. Natur. Hist. Ser., 10: 174—177.
- Berland, B. 1970. on the morphology of head in four species of Cucullanida (Nematoda). Sarsia, 43: 15—63.
- Bilqees, F.M., 2007. Important helminth parasites of Pakistan. Fed.Urdu Univ. Sci. Techn. Karachi. pp. 1-439.
- Bilqees, F. M., Khanum, Z., & Jehan, Q. 1971. Marine fish Nematodes of West Pakistan I. Description of seven new species of Karachi coast J. Sci. Karachi, 1 (1) : 175-184.
- Bilqees, F.M., Akhtar, Y., Haseeb, M.F. & Khalil, B. 2005. A new nematode *Cucullanus mujibi* n.sp.(cucullanidae Cobbold, 1864) from the fish *Arius serratus*(Day) of Karachi coast. Proc. Parasitol; 39:79-92.
- Bilqees, F.M. and Fatima, H. (1980a). Marine fish nematodes of Pakistan XII. *Cucullanus quadrii*, new species (Cucullanidae) from *Arius serratus* (Day.) of Karachi coast. Pak. J.Zool., 12:27-31.
- Caspeta-Mandujano, J.M., Moravec, F. & Aguilar-

- Aguilar, R. 2000. *Cucullanus mexicanus* sp.n.(Nematoda:Cucullanidae) from the intestine of the freshwater catfish *Rhamdia guatemalensis*(Pimelodidae) in Mexico *Helminthologia* 37: 215-217.
- Chandler, A. C. 1935. Parasites of fishes in Galveston Bay. *Proc. U.S. natn. Mus.*, 83: 123-157.
- David Gonzalalez-Solis., Vielka, M. Tuz-Paredes and Miguel, A. Quintal-Loria. 2007. *Cucullanus pargi* sp.n.(Nematoda: Cucullanidae) from the grey snapper *Lutjanus griseus* off the southern coast of Quintana Roo, Mexico, *Folia. Parasitol.* 54: 220-224.
- Fortes, E., Hoffmann, R.P. and Sarmiento, J.M. 1993a. Nova especie de nematodeo *Cucullanus*, *C. riograndensis* sp.n., de *Pimelodus maculatus* Laeepede, 1803(Pisces), do Lago do Guiba, Rio Grande do Sul, Brasil. *Rev. Bras. Med. Vet.*, 15: 79-82.
- Fortes, E., Hoffmann, R.P. and Sarmiento, J.M. 1993b. *Cucullanus fabrigasi* sp. N. nematoda parasita do intestino do peixe *Pimelodus maculatus* (Lacepede, 1803) do Rio Guaiba, Poroto Alegre, Rio Grande do Sul, Brasil. *Arq. Fac. Veter. UFRGS. No.21:* 71-76.
- Gendre, E. 1927. *Parasitologia mauritanica*. Nematodes parasites des poissons de la cote de Mauritanie. *Bull. Com. Etud. Hist. Scient. Afr. Occid. Fr.*, 10: 258-274.
- Gupta, S. P.& Gupta, R. C. 1979. On some nematode parasites of marine fishes. *Indian J. Helminth.*, 29 (1977): 104-112.
- Gupta P.C., Masoodi B.A. 1982. Three new and one known piscine nematodes from Kanpur. *Kanpur Univ. Res. J. (Sci.)* 3:57-70.
- Gupta, S.P. and Naqvi, N.H. 1983. Nematode parasites of fishes. VIII. On two new species of the genus *Indocucullanus* Ali, 1956 from fishes. *Indian. J.Helminth.* 34(1): 78-85.
- Gupta, S.P. and Srivastava, A.B. 1984a. On three new nematode parasites (Nematoda: Cucullanidae) from Indian fishes. *Acta Parasitologica Polonica*, 29(10): 77-84.
- Ingrid, D.V., Tomas, T.J. and Haydgc, S.N. 2002. *Cucullanus marplatensis* sp. Nov. (Nematoda: Susullanidae) parasitizing *Odontesthes argentinensis* (Valenciennes, 1835) (Pisces, Atherinidae) from Argentinean water. *Acta Parasit.*, 47(1): 41-46.
- Kalyankar, S. D. 1971. On some nematodes from Indian with the description of a new species (Ascaridodidea: Stomachidae). *Riv. Parasitol.*, 33: 203—208.
- Karve, J. N. 1952. Some parasitic nematode of fishes. III. *J. Univ. Bombay*, 21 (3) : 1-14.
- Køie M .2000. The life-cycle of the flatfish nematode *Cucullanus heterochrous*. *J Helminthol* 74:323-328.
- Lanfranchi, L. Ana ; Juan T. T. and Norma, H .S.2004. *Cucullanus bonaerensis* n. sp. (Nematoda: Cucullanidae) parasitizing *Urophycis brasiliensis* (Pisces: Phycidae) from Argentinean waters. *J. parasitol.* : 90(4) 808-812.
- Lane, C. 1916. The genus *Dacnitis* Duj, 1845. *Indian J. Med. Res.*, 4: 93-104.
- Lakshmi, B. B. 2000. A new parasite of the genus *Cucullanus*(Nematoda: Cucullanidae) from the fish *Thalassinus*(Ruppel.) *Biol. Chil. Parasitol.*, 55: 3-4.
- Lebre, C. and Peter, A. J. 1984. *Cucullanus campanae* n. sp. (Cucullanidae, Nematoda), a parasite of *Solea vulgaris vulgaris*,. *Bull. Du Mus.d’Histoire Naturelle*, 6: 999-1005.
- Moravec, F. 1979. Observation on the development of *Cucullanus*(*Truttaedacnitis*) *truttiae* (Fabricius,

- 1794) (Nematoda:Cucullanidae) *Folia Parasitol.* 26: 295-307.
- Moravec, F. 1983. Remarkable development of *Cucullanus truttae*, a parasite of trout in Czechoslovakia *Ziva.* 2: 63-64.
- Moravec, F., Kohn, A., Fernandes, B.M.M. 1993. Nematode parasites of fishes of the Parana River, Brazil. Part 2. Scuratoidea, Ascaridoidea, Habronematoidea and Acuarioidea: *Folia Parasitol.*, 40: 115-134.
- Moravec, F.; Lorber, J. and Konecny, R. 2008. *Cucullanus maldivensis* n. sp. (Nematoda: Cucullanidae) and some other adult nematodes from marine fishes off the Maldiv Islands. *Syst. Parasitol.* 70(1):61-9.
- Moravec, F., Sasal, P., Wurtz, J. & Taraschewski, H. 2005. *Cucullanus oceaniensis* sp.n. (Nematoda: Cucullanidae) from Pacific eel (*Anguilla* spp.) *Folia Parasitologica* 52:343-348.
- Petter, A.J. 1974. Essai de classification de la famille des Cucullanidae.
- Bull. Mus. Natl. Hist. Nat. Paris, Series 3, Zoologie,* 177: 1469-1491.
- Rasheed, S. 1968. The nematodes of the genus *Cucullanus* Muller, 1777, from the marine fish of Karachi coast. *An. Esc. Nac. Cienc. Biol. Mex.*, 15: 23-59.
- Rehana, R. & Bilquees, F. M. 1986. Redescription of *Cucullanus annulatus* Rehana and Bilquees 1976 as *Cucullanus pseudoannulatus*, new species from *Mystus cavasius* (Ham.) of Kalri lake, Sind, Pakistan. *J.zool.*, 18: 345-349.
- Sood, 1989. *Fish Nematodes from South Asia.* Kalyani Publ. New Delhi, pp704.
- Soota, T. D. 1983. Studies on nematode parasites of Indian vertebrates I. Records of the Zoological survey of India, occasional paper no.54 *Fishes.* 1-352.
- Soota, T. D. and Chaturvedi, Y. 1971. On five new nematodes from vertebrates. *Zool. Anz.*, 187: 310-317.
- Soota, T. D. & Dey Sarkar, S. R. 1980. On three species of the nematode genus *Cucullanus* Mueller, 1777, and a note on *Lappetascaris lutjani* Rasheed, 1965, from Ind. Mar. fish. Rec. *Zool. Surv. India.*, 76:1-6.
- Srivastava, A. B. & Gupta, S. P. 1976. Nematode parasites of fishes 2. On two new species of *Indocucullanus* Ali, 1956. *Japan J. Parasit.*, 26:46-48.
- Timi, J.T. and Lanfranchi, A.L. 2006. A new species of *Cucullanus* (Nematoda: Cucullanidae) parasitizing Conger *orbignianus* (Pisces: Congridae) from Argentinean waters. *J. Parasitol.* 92: 151-154.
- Vicente, J.J. and Santos, E. Dos. 1974. Helminth of fish from the north Fluminense littoral. *Mem. Inst. Oswaldo Cruz.*, 27: 173-180.
- Vicente, J.J. and Fernando, B.M.M. 1973. On a new nematode of the genus *Cucullanus* Mueller, 1777, parasite of *Norurado* fish (Nematoda: Camallanidae). *Atas de Sociedade Biol. Rio. Janeiro.*, 17: 31-33.
- Wang, P.Q. 1984. Descriptions of three new species and a list of parasitic nematodes from vertebrates in Fujian province. *Wuyi Sci. J.*, 4: 113-132.
- Yamaguti, S. 1935. Studies of helminth fauna of Japan. Pt. 9. Nematodes of fishes 1. *Japan. J. Zool.*, 9: 337-386.
- Yamaguti, S. 1941. Studies on the helminth fauna of Japan. pt. 33. Nematodes of fishes. II. *Jap. J. Zool.*, 9(3): 343-396.

Yamaguti, S. 1954. Parasitic worms mainly from Celebes. Pt. 9. Nematodes of fishes. Acta. Med. Okoyama, 9(1): 122-133.

Yamaguti, S. 1961. Systema Helminthum. Vol. III. The nematodes of vertebrates, pts. I & 2, Intersc. Publishers: 1-1261.

Zaidi, D. A. and Khan, D. 1975. Nematode parasites from fishes of Pakistan. Pakistan. J. Zool; 7: 51-73.