

Prevalence of Cestode Parasites of *Gallus Domesticus* in Karachi, Sindh, Pakistan

Rubab Malik*, Nasira Khatoon

Department of Zoology, University of Karachi, Karachi-75270, Pakistan.

ABSTRACT

Background: Poultry production in Pakistan is growing remarkably and contributing significantly to the economy. However, disease outbreaks are the most serious threat to poultry productivity. Among other poultry diseases which cause huge damage to poultry, parasitic diseases are also included, but no significant work has been done on this area of concern in Pakistan.

Objective: This study is carried out to evaluate the prevalence rate of one of the major helminth parasite groups, Cestodes, and their effects on the health of chickens in the districts of Korangi, Karachi, Sindh.

Methodology: Between February 2018 and January 2019, 76 chickens were examined for Cestodes infection. Cestodes were extracted from the gut and fixed in F.A.A. fixative for 24 hours with a little cover slip pressure. The cestodes were then thoroughly cleaned with 70% ethanol, stained with Mayer's carmalum, dehydrated in progressively stronger alcohol, clarified with clove oil, rinsed with xylene, and permanently mounted in Canada balsam.

Results: The overall prevalence was 52.6%. Parasite burden is high in males (63.8%) compared to females (45.6%). Six species of Cestodes parasites were identified from the intestine of the infected chickens. *Raillietina tetragona*, *R. cesticillus*, *R. echinobothridia*, *R. ransomi*, *Choanotaenia infundibulum*, and *Raillietina* sp. are the Cestode species that have been found. Their prevalence was 42.1%, 34.2%, 26.3%, 22.2%, 15.7%, and 7.8%, respectively. The prevalence rate also varies from season to season. Spring had the highest rate of infection at 60%, summer was next at 58%, and autumn and winter both had a rate of 40%.

Conclusion: The study concludes that heavy infestation has a negative impact on the physical health of chickens and points to the need to control Cestodes parasite infection in order to reduce mortality and increase poultry yield.

Keywords

Poultry, Prevalence, Cestodes parasites, chickens, Karachi.

*Address of Correspondence

rubabmalik943@gmail.com

Article info.

Received: December 12, 2021

Accepted: November 01, 2022

Cite this article: Malik R, Khatoon N. Prevalence of Cestode Parasites of *Gallus Domesticus* in Karachi, Sindh, Pakistan. 2022; 13(2):157-162.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium provided the original work is properly cited.

INTRODUCTION

In Pakistan, the poultry industry is a big part of the economy because it makes eggs and meat, which are the cheapest sources of protein. It contributes 29% of the total meat production of the country, 5.8% of the agriculture sector, and 1.3% of the overall GDP. It has shown excellent growth and become a source of 1.6 million people's employment. Among other poultry diseases which cause huge damage to poultry, parasitic diseases are also

included, but no significant work has been done on this area of concern in Pakistan. The common gastrointestinal helminth parasites cause negative effects on the health of chickens, which then leads to considerable damage and great economic losses to the poultry industry due to malnutrition, decreased food conversion ratio, weight loss, lower egg production, and death in young birds¹. Parasites also make the flocks less resistant to diseases and make diseases worse^{2,3}. Chicken cestodiasis not only causes loss of body weight but also may cause several problems,

such as enteritis, loss of blood, loss of production, nervous manifestations, and death. Parasitic infection causes villous atrophy, catarrhal enteritis, granuloma formation in the duodenum, desquamation of villi and submucosal gland congestion, an inflammatory response, and vacuolation of epithelial cells⁴. However, poultry production in Pakistan is growing remarkably and contributing significantly to its economy. However, disease outbreaks pose the greatest threat to productivity. Among other diseases that cause huge damage to poultry, parasitic diseases are also included, but no significant work has been done on this area of concern in Pakistan. It is necessary to investigate the rate of infection of various cestode parasites in chickens. The current research aims to study the prevalence and intensity of infection of cestode parasites of *Gallus domesticus* and to study the difference in rate of parasitic infection in different seasons and months to analyse the effect of seasonal variation on parasitic infection.

MATERIALS AND METHODS

The viscera of 76 freshly slaughtered chickens have been collected from common shops located in Korangi, Karachi, from February 2017 to January 2018. Samples were subjected to necroscopic and parasitological examination. Cestodes were collected from the gastrointestinal tract and preserved in Acetic acid Formalin Alcohol solution (AFA) for 8 to 12 hours. Then it is kept in 70% alcohol overnight and stained with Alum carmine for 3-5 minutes and is continuously checked to prevent overstrain. Then it is dehydrated through 70%, 90% and 100% alcohol for 20-30

minutes for each grade. It is then bathed in clove oil for 1-2 minutes and washed in xylol for 2-3 minutes. Finally, permanent slides were prepared by using Canada balsam. Cestodes were identified under a light microscope with 10x magnification according to their morphological characteristics⁵. However, Pearson's coefficient of correlation 'r' was used to examine the relationship among seasons and infection prevalence. At the 5% threshold of significance ($P \leq 0.05$), statistical significance was recognised. SPSS version 20 was used to analyse the data.

RESULTS AND DISCUSSION

The author intends to investigate the prevalence of cestode parasite infection in the Korangi Creek area. The sampling area is selected due to the author's residence in the area, which provides easy access to get samples. The study found that, out of 76 chickens, 40 had different species of Cestodes (Table 1). Six species belonging to three genera were found in the digestive tract of chickens. The overall prevalence of cestodes was 52.6%. Parasite burden is high in males (63.8%) as compared to females (45.6%) (Table 2).

Six species of Cestodes parasites were found in the intestines of infected chickens. *Raillietina tetragona*, *R. cesticillus*, *R. echinobothridia*, *R. ransomi*, *Choanotaenia infundibulum*, and *Raillietina* species are the species of Cestodes that have been found. Their prevalence was 42.1%, 34.2%, 26.3%, 22.2%, 15.7%, and 7.8%, respectively (Table 3).

Table 1. Overall prevalence of Cestodes infection.

Helminthes group	Host Examined	Host Infected	Prevalence %
Cestodes	76	40	52.6

Table 2. Prevalence of Cestodes in relation to host's sex.

Sex of host	Host examined	Host infected	Prevalence (%)
Male	30	19	63.8
Female	46	21	45.6

Table 3. Prevalence of Cestodes parasites.

Genus	Host examined	Host infected	Prevalence (%)
<i>Raillietina tetragona</i>	76	32	42.1
<i>Raillietina cesticillus</i>	76	26	34.2
<i>Raillietina echinobothridia</i>	76	20	26.3
<i>Raillietina ransomi</i>	76	17	22.2
<i>Choanotaenia infundibulum</i>	76	12	15.7
<i>Raillietina</i> sp.	76	6	7.8

Table 4. Seasonal incidence of recorded Cestodes' Species.

Season	Host Examined	Host Infected	Prevalence %
Spring	15	9	60.0
Summer	36	22	58.3
Autumn	15	6	40.0
Winter	10	3	40.0

Raillietina tetragona was reported in chicken 48.3%⁶ and 51.66%⁷ from Pakistan; 14.5%⁸ from Western Cameroon; 16.2%⁹ from Egypt; and 2.0%¹⁰ from Nigeria; 7.4%¹¹ from Egypt; 51.42%¹²; and 9.16%¹³ from India; 57.5%¹⁴ from Thailand; 68.75%¹⁵ from Algeria; 19.68%¹⁶ from Bangladesh; and 65.94%¹⁷ from Ethiopia

R. echinobothrida reported from the chicken with prevalence of 19.2%⁶ from Pakistan, 10.4%¹⁸ from Iran, 48.3%¹⁴ from Northern Thailand, 25.3%¹⁹ from Nigeria, from Brazil²⁰, 91.9%²¹ and 6.6%¹¹ from Egypt, 39.86%¹⁷ from Ethiopia, 83.3%¹⁵ from Algeria, 37.83%²² from Phitsanulok province, 2.3%²³ from Trinidad, 50%²⁴ from Bangladesh and 46.3%⁹ from Tanzania while the same specie has been reported in pigeon 61.76%²⁵ from Greece and 32%²⁶ from Libya.

R. cesticillus was reported from birds with prevalence of 56.6%²⁴ from Bangladesh, 55.07%¹⁷ from Ethiopia, 83.5%²⁷ from Pakistan, 23.22%²⁸ from India, 29.1%¹⁵ from Algeria and 21.3%¹⁹ from Nigeria, 0.9%²³ from Trinidad, 5.83%⁷ and 0.8%⁶ from Pakistan, 12.5%¹⁴ from Thailand and, also reported from Brazil²⁰.

Choanotaenia infundibulum was reported from Hyderabad, Pakistan, with a prevalence of 89.5%²⁷, while *Choanotaenia* sp. with prevalence of 20% reported from the same host in Quetta, Pakistan⁶. Outside of Pakistan, the same species has been reported from Zimbabwe²⁹, from Trinidad²³, from Tanzania⁹, from Algeria¹⁵, from Japan³⁰ and from Brazil³¹.

The difference in prevalence might be due to the variation in the environmental conditions of the area³², including high temperature, humidity or dryness, annual rainfall, and the availability of intermediate hosts³³, it could also be due to sharing the same habitat with migratory birds, which increases the chance of harbouring parasites³⁴. Similar attribution is also given that environmental alterations, especially increasing temperature, may have affected the occurrence of helminth infection¹⁴, and another author stated that more than one million birds migrate from Central Asia to Pakistan during winters. During migration, they also carry parasites, which result in new host and locality records³⁵.

Moreover, the study found that the prevalence rate also varies from season to season. The highest infection has been observed in spring at 60%, followed by summer at 58.3%, and in autumn and winter, the same infection has been observed at 40% (Table 4). This suggested that there was not enough variation in the prevalence and intensity of infection in different seasons. The results suggest that high temperatures increase the susceptibility of the hosts to parasitism⁶. Although parasites were discovered to be prevalent throughout the year, with a higher prevalence and parasitic load, they were found to be most prevalent in the summer season.

Furthermore, multiple infections of helminth parasites have also been found in many chickens. Similarly, a high intensity of parasites has also been observed. The intestine

was fully blocked with the worm burden. Grossly, the intestine showed a heavy infestation of Cestodes, due to which the mucosal lining of the intestine was damaged. Chronic catarrhal enteritis, hemorrhagic enteritis, and

nodular enteritis have been observed. The intestinal wall was thickened and inflamed with hemorrhagic necrotic spots.

Table 5. Seasonal incidence of each specie of recorded Cestodes.

Season	<i>R.tetragona</i>		<i>R.cesticillus</i>		<i>R.echinobothrida</i>		<i>R.ransomi</i>		<i>Choanotaenia Infundibulum</i>		<i>Raillietina sp.</i>		P-Value
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
Spring	8	88.8	6	66.6	4	44.4	2	22.2	2	22.2	0	0	0.028
Summer	17	77.2	15	68.1	13	59	11	50	8	36.3	4	18.1	0.002
Autumn	5	83.3	4	66.6	3	50	2	33.3	2	33.3	1	16.6	0.005
Winter	2	66.6	1	0	0	0	2	66.6	0	0	1	33.3	0.041

Table 6. Monthly incidence of different types of recorded Cestodes' species.

Month	<i>R.tetragona</i>		<i>R.cesticillus</i>		<i>R.echinobothridia</i>		<i>R.ransomi</i>		<i>Choanotaenia infundibulum</i>		<i>Raillietina sp.</i>		P-Value
	No	%	No	%	No	%	No	%	No	%	No	%	
Feb	3	100	2	66.6	2	66.6	1	33.3	1	33.3	0	0	0.017
Mar	2	100	2	100	1	50	0	0	0	0	0	0	0.093
Apr	3	75	2	50	1	25	1	25	1	25	0	0	0.025
May	7	77.7	7	77.7	6	66.6	5	55.5	3	33.3	2	22.2	0.002
Jun	3	75	3	75	2	50	2	50	2	50	1	25	0.001
Jul	7	77.7	5	55.5	5	55.5	4	44.4	3	33.3	1	11.1	0.004
Aug	1	100	0	0	1	100	1	100	1	100	1	100	0.004
Sep	2	100	2	100	1	50	1	50	1	50	0	0	0.013
Oct	2	66.6	2	66.6	1	33.3	0	0	0	0	0	0	0.093
Nov	1	50	1	50	0	0	2	100	0	0	0	0	0.102
Dec	0	0	0	0	0	0	0	0	0	0	0	0	0
Jan	1	100	0	0	0	0	0	0	0	0	1	100	0.175

CONCLUSION

The present study concludes that the prevalence of parasitic infection by cestodes is very high, resulting in the mortality of chickens. The high prevalence of infection leads to invisible production and economic losses. Rearing area, season, availability, and contact with the intermediate host also affect the prevalence and intensity of infection. Based on the result, the following recommendations are forwarded: As the cestodes parasites have a sub-clinical occurrence, studies that focus on these ideas should be conducted. The public, especially those who are related to poultry farming, should be aware of the occurrence and economic significance of gastrointestinal parasites. Elimination of intermediate hosts in the rearing area and extensive early-season larval control have been recommended.

ETHICAL APPROVAL

Not required.

CONFLICTS OF INTEREST

No conflict of interest.

FUNDING SOURCE

None.

ACKNOWLEDGEMENTS

I am thankful to my supervisor for her guidance and support.

LIST OF ABBREVIATIONS

AFA Acetic acid Formalin Alcohol
GDP Gross Domestic Product

REFERENCES

1. Asumang P, Akoto Delali J, Wiafe F, Kamil Z, Iddrisu Balali G, Afua Dela Gobe V, et al. Prevalence of Gastrointestinal Parasites in Local and Exotic Breeds of Chickens in Pankrono–Kumasi, Ghana. *Journal of parasitology research*. 2019 Sep 2;2019.
2. Lamy E, van Harten S, Sales-Baptista E, Guerra MM, de Almeida AM. Factors influencing livestock productivity. Environmental stress and amelioration in livestock production. 2012;19-51.
3. Kaur H, Katoch A. Prevalence, site and tissue preference of myxozoan parasites infecting gills of cultured fish in Punjab (India). *Dis Aquat Organ*. 2016;118(2):129-37.
4. Kurkure NV, Kolte SW, Ganorkar AG, Bhandarkar AG. *Raillietina cesticillus* in pigeon (*Columbia livia*). *Indian Vet J*. 1998;75(9):835-6.
5. Yamaguti S. *Systema helminthum*. Vol. II. The cestodes of vertebrates. *Systema helminthum*. Vol. II. The cestodes of vertebrates. 1959.
6. Ahmed S, Babar S, Fareed SK, Kakar MA, Saadullah JA. Helminthosis of rural poultry in Quetta, Pakistan. *Eurasian J Vet Sci*. 2013;29(2):103-5.
7. Tasawar Z, Aziz F, Akhtar M. Prevalence of cestode parasites of domestic fowl (*Gallus gallus domesticus*). *Pak Vet J*. 1999;19(3):142-4.
8. Mpoame M, Agbede G. The gastro-intestinal helminth infections of domestic fowl in Dschang, western Cameroon. *Rev Elev Med Vet Pays Trop*. 1995;48(2):147-51.
9. El-Dakhly KM, El-Seify MA, Mohammed ES, Elshahawy IS, Fawy SA, Omar MA. Prevalence and distribution pattern of intestinal helminths in chicken and pigeons in Aswan, Upper Egypt. *Tropical Animal Health and Production*. 2019 Mar 12;51:713-8.
10. Ola-Fadunsin SD, Uwabujo PI, Sanda IM, Ganiyu IA, Hussain K, Rabiun M, Elelu N, Alayande MO. Gastrointestinal helminths of intensively managed poultry in Kwara Central, Kwara State, Nigeria: Its diversity, prevalence, intensity, and risk factors. *Vet World*. 2019;12(3):389.
11. Nagwa EA, Loubna MA, El-Madawy RS, Toulou EI. Studies on helminths of poultry in Gharbia governorate. *Benha Vet Med J*. 2013;25(2):139-44.
12. Ayshia A, Wani SA. Endohelminth parasites of domestic fowl (*Gallus domesticus*) in Doda district of Jammu & Kashmir State, *Indian Vet J*. 2015;13(1):39-42.
13. Singh H, Nama P. Incidence of endohelminth parasites in the alimentary canal of domestic fowl (*Gallus domesticus*), butchered at Pipar city, Jodhpur. *Int J Res Anal Rev*. 2018;5:130-3.
14. Butboonchoo P, Wongsawad C. Occurrence and HAT-RAPD analysis of gastrointestinal helminths in domestic chickens (*Gallus gallus domesticus*) in Phayao province, northern Thailand. *Saudi J Biol Sci*. 2017;24(1):30-5.
15. Ilyes M, Ahmed B. Cestode parasites of free-range chickens (*Gallus gallus domesticus*) in the north-eastern of Algeria. *Int J Poult Sci*. 2013;12(11):681-4.
16. Alam MN, Mostofa M, Khan MA, Alim MA, Rahman AK, Trisha AA. Prevalence of gastrointestinal helminth

- infections in indigenous chickens of selected areas of Barisal district, Bangladesh. *Bangladesh J Vet Med.* 2014;12(2):135-9.
17. Berhe M, Mekibib B, Bsrat A, Atsbaha G. Gastrointestinal helminth parasites of chicken under different management system in Mekelle town, Tigray region, Ethiopia. *J Vet Med.* 2019;2019.
 18. Ali Mohammad B, Mani D, Minoo S. Study of parasitic infection and its histological changes in a bird. 2015.
 19. Adang KL, Asher R, Abba R. Gastro-intestinal helminths of domestic chickens *Gallus gallus domestica* and ducks *Anas platyrhynchos* slaughtered at Gombe main market, Gombe State, Nigeria. *Int J Poult Sci.* 2014;8(2):32-40.
 20. da Silva GS, Romera DM, Fonseca LE, Meireles MV. Helminthic parasites of chickens (*Gallus domesticus*) in different regions of São Paulo State, Brazil. *Rev Bras Cienc Avic.* 2016;163-8.
 21. Shahin AM, Lebdah MA, Abu-Elkheir SA, Elmeligy MM. Prevalence of chicken cestodiasis in Egypt. *NY Sci J.* 2011;4(9):21-9.
 22. Phalee W, Phalee A. Cestode Diversity of Free-Range Domestic Chickens (*Gallus gallus domesticus*) in Phitsanulok Province. *Burapha Sci J.* 2017;22(1):271-8.
 23. Baboolal V, Suratsingh V, Gyan L, Brown G, Offiah NV, Adesiyun AA, Basu AK. The prevalence of intestinal helminths in broiler chickens in Trinidad. *Vet Arh.* 2012;82(6):591-7.
 24. Begum A, Mukutmoni M, Akter F. Parasite Diversity in Domestic Duck on *Anas Platyrhynchos Domesticus* from Munshiganj, Dhaka. *Bangladesh J Zool.* 2018;47(1):121-8.
 25. Diakou A, Ptochos S, Papadopoulos E. Cestode fauna of feral pigeons in Thessaloniki; Northern Greece. *Helminthologia.* 2013;50:39-42.
 26. Alkharigy FA, El Naas AS, Maghrbi AA. Survey of parasites in domestic pigeons (*Columba livia*) in Tripoli, Libya. *Vet J.* 2018;8(4):360-6.
 27. Butt Z, Shaikh AA, Memon SA, Narejo NT, Jalbani S. 03. Pathology of *Choanotenia infundibulum* in the intestine of free range chickens from Hyderabad, Sindh, Pakistan. *Pure and Applied Biology (PAB).* 2021 Oct 18;4(2):158-63.
 28. Salam ST, Mir MS, Khan AR. The prevalence and pathology of *Raillietina cesticillus* in indigenous chicken (*Gallus gallus domesticus*) in the temperate Himalayan region of Kashmir-short communication. *Vet Arh.* 2010;80(2):323-8.
 29. Percy J, Pias M, Enetia BD, Lucia T. Seasonality of parasitism in free range chickens from a selected ward of a rural district in Zimbabwe. *Afr J Agric Res.* 2012;7(25):3626-31.
 30. Gamra OW, Antia RE, Falohun OO. Intestinal cestodes of poultry *Raillietina echinobothrida* and *Choanotaenia Infundibulum* infection in a commercial Japanese quail (*Coturnix coturnix japonica*) farm in Apomu, Osun State, Nigeria. *Sci J Zool.* 2015;4(4):20-5.
 31. Monte GL, Cavalcante DG, Oliveira J. Parasitic profiling of Japanese quails (*Coturnix japonica*) on two farms with conventional production system in the Amazon region. *Pesqui Vet Bras.* 2018;38:847-51.
 32. Bennema SC, Vercruyse J, Morgan E, Stafford K, Höglund J, Demeler J, von Samson-Himmelstjerna G, Charlier J. Epidemiology and risk factors for exposure to gastrointestinal nematodes in dairy herds in northwestern Europe. *Veterinary Parasitology.* 2010;173(3-4):247-54.
 33. Charlier J, Thamsborg SM, Bartley DJ, Skuce PJ, Kenyon F, Geurden T, Hoste H, Williams AR, Sotiraki S, Höglund J, Chartier C. Mind the gaps in research on the control of gastrointestinal nematodes of farmed ruminants and pigs. *Transbound Emerg Dis.* 2018;65:217-34.
 34. Besier RB, Kahn LP, Sargison ND, Van Wyk JA. The pathophysiology, ecology and epidemiology of *Haemonchus contortus* infection in small ruminants. *Adv Parasitol.* 2016;93:95-143.
 35. Birmani NA, Dharejo AM, Khan MM. *Catantropis* sp. (Trematoda: Notocotylidae) from the Black Coot, *Fulica atra* Linnaeus, 1758 (Gruiformes: Rallidae) in Sindh Province of Pakistan. *JAPS, J of Animal and Plant Sci.* 2011;21(4):872-3.